Using Older Observations to Improve Ephemerides

E M Standish(JPL/Caltech)

There is a wild variety of older planetary observations, each wit Irits own set of circumstances, techniques and characters. Galileo, in 1012, observed Neptune and knew there was something unusual shout that observation: he almost, . . . Lalande, in 1795, also observed Neptune and also noted something unusual about his observations; he, too, almost . . . Both of these Neptune observations are useful in refining our presently determined orbit for Neptune.

There are other observations, however, which cannot be trusted, and the reasons for these sometimes bor der upon the sinister. Two sets, in particular, are examined here: the too-good- to-be-true transit observations of Abram Robertson in 1S1 1 and the suspicious eclipse measurements of Samuel Williams in 1780.

Abstract submitted for AAS [11A])] meeting

O

Date submitted: October 9, 1995 Electronic form version 1.6

[American Astronomical Society Abstract Form]

AAS Category 26

Running # π

Session 0.00

Oral preferred X

Dissertation abstract [-1

Paper presented by E M Standish

JPL/CalTech 301-150 Pasadena, CA 91109 818-354-3959

ems@smyles.jpl.nasa.gov

Special instructions:

Abstract fee of \$40 will be paid by PO

Purchase order billing data: 1'.(). #tbaupon receipt of invoice

JPL 301-125

Pasadena, CA 91109 attn: Bob Klotz